Neuroscience Letters, 158 (1993) 237–239
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Adachi, A., see Kobashi, M., 47
Adams, M.E., see Geer, J.J., 97
Albrecht, J., see Matyja, E., 155
Amadeo, A., see Spreafico, R., 232
Andou, Y., see Zhang, L., 105
Aou, S., Ma, J. and Hori, T., Hypothalamic involvement in stress-induced hypocalcemia in rats, 197
Arisawa, M., see Kuwahara, T., 1

Bandou, Y., see Kusakabe, T., 59
Barker, J.L., see Vautrin, J., 125
Bernardi, G., see Sancesario, G., 101
Block, J.J., see O'Donoghue, D.L., 143
Bowery, N.G., see Malcangio, M., 189
Brookes, S.J.H., see Song, Z.-M., 221

Całka, J., Majewski, M., Kaleczyc, J. and Łakomy, M., Immunocytochemical demonstration of neuropeptide Y and luteinizing hormone-releasing hormone-immunoreactive structures in the organum vasculosum laminae terminalis of juvenile gilts, 21
Calvo, P., see Negro, M., 83
Casanova, E., see Negro, M., 83
Casas, R., see Ferrer, I., 163
Castañeda, E., see Pellis, S.M., 13
Chang, C.C., see Hong, S.J., 25
Chiba, T., see Tanaka, K., 67
Chinchetru, M.A., see Negro, M., 83
Ciriello, J., see Hochstenbach, S.L., 225
Connelly, C.D., see Raffa, R.B., 87
Costa, M., see Song, Z.-M., 221
Craig, C.G. and White, T.D., NMDA-evoked adenosine release from rat cortex does not require the intermediate formation of nitric oxide, 167
Crusio, W.E., see Nek, N., 75

DasGupta, B.R., see Schengrund, C.-L., 159
De Biasi, S., see Spreafico, R., 232
De Blas, A.L., see Spreafico, R., 232
Demura, T., see Matsuura, T., 5
Dooley, D.J., see Geer, J.J., 97
Doyu, M., see Yamamoto, M., 39
Droge, M.H. and Tao, Y., Glycine effects on in vitro motor pattern generation in mouse spinal cord, 139
Dudel, J., Franke, C. and Luboldt, W., Reaction scheme for the glutamate-ergic, quisqualate type, completely desensitizing channels on crayfish muscle, 177
Dun, N.J., Dun, S.L., Wu, S.Y. and Förstermann, U., Nitric oxide synthase immunoreactivity in rat superior cervical ganglia and adrenal glands, 51
Dun, S.L., see Dun, N.J., 51

Eggeman, K.T., Pal, R., Walsh, J., Kumar, K.N. and Michaelis, E.K., Immunochemical and immunohistochemical characterization of a synaptic membrane protein that binds the competitive antagonists of NMDA receptors, 173
Emelyanov, N.A., see Mokrushin, A.A., 16
Endo, T., see Go, M., 135
Erneux, C., see Go, M., 135
Esumi, H., see Tanaka, K., 67
Esumi, H., see Ohta, A., 33

Fernández-López, A., see Negro, M., 83
Ferrer, I., Zújar, M.J., Rivera, R., Soria, M., Vidal, A. and Casas, R., Parvalbumin-immunoreactive dystrophic neurites and aberrant sprouts in the cerebral cortex of patients with Alzheimer's disease, 163
Förstermann, U., see Dun, N.J., 51
Franke, C., see Dudel, J., 177
Frotscher, M., see Nek, N., 75
Fusco, F., see Sancesario, G., 101

Gajendiran, M., see Seth, P., 217
Ganguly, D.K., see Seth, P., 217
Gean, P.-W. and Lin, J.-H., D-2-Amino-5-phosphonovaleate blocks induction of long-term depression of the NMDA receptor-mediated synaptic component in rat hippocampus, 170
Geer, J.J., Dooley, D.J. and Adams, M.E., K^+-stimulated $^{45}Ca^{2+}$ flux into rat neocortical mini-slices is blocked by ω-Aga-IVA and the dual Na^+/Ca^{2+} channel blockers lidoflazine and flunarizine, 97
Giacobini, E., see Messamore, E., 205
Ginsberg, S.D., Hof, P.R., McKinney, W.T. and Morrison, J.H., The noradrenergic innervation density of the monkey paraventricular nucleus is not altered by early social deprivation, 130
Go, M., Uchida, T., Takazawa, K., Endo, T., Erneux, C., Mailleux, P. and Onaya, T., Inositol 1,4,5-trisphosphate 3-kinase highest levels in the dendritic spines of cerebellar Purkinje cells and hippocampal CA1 pyramidal cells. A pre- and post-embedding immunoelectron microscopic study, 135
Gong, X.-W., see Hashimoto, M., 29
Green, P.G., Luo, J., Hammond, E.R. and Levine, J.D., Trypsin enhances sympathetic neuron-dependent plasma extravasation in the rat knee joint, 117

Hamai, Y., see Ohta, A., 33
Hammond, E.R., see Green, P.G., 117
Hashiguti, H., see Nakahara, D., 44
Hashimoto, M., Gong, X.-W., Izaki, Y., Iriki, M. and Hibino, H., 1-Oleoyl-2-docosahexaenoyl phosphatidylcholine increased paradoxical sleep in F344 rats, 29
Hayashi, H., see Toda, T., 36
Henselmans, J.M.L., Heyna, M.H. and Stoof, J.C., Regional differences in the dopaminergic regulation of cyclic AMP formation within the rat nucleus accumbens: comparison with the striatal complex of the lizard *Gekko gecko*, 79
Heyna, M.H., see Henselmans, J.M.L., 79
Hibino, H., see Hashimoto, M., 29
Hirai, H., see Ikeuchi, Y., 63
Hochstenbach, S.L., Solano-Flores, L.P. and Ciriello, J., Fos induction in brainstem neurons by intravenous hypertonic saline in the conscious rat, 225
Hof, P.R., see Ginsberg, S.D., 130
Hong, S.J. and Chang, C.C., O_2 occlusion and cyanide induced immediate relaxation and contraction of murine skeletal muscle, 25
Hori, H., see Kusakabe, T., 59
Hori, T., see Aou, S., 197

Ichikawa, H., see Kobashi, M., 47
Iida, S., see Ohta, A., 33
Ikeda, J., Ochiai, K., Morita, I. and Murota, S.-i., Endogenous nitric

oxide blocks calcium influx induced by glutamate in neurons containing NADPH diaphorase, 193
Ikeuchi, Y., Hirai, H., Okada, Y., Mio, T. and Matsuda, T., Excitatory and inhibitory effects of toluene on neural activity in guinea pig hippocampal slices, 63
Iriki, M., see Hashimoto, M., 29
Izaki, Y., see Hashimoto, M., 29

Kaleczyc, J., see Całka, J., 21
Kamei, J., Kawashima, N., Suzuki, T., Misawa, M. and Kasuya, Y., The effects of cyclosporine on morphine-induced antinociception in diabetic mice, 213
Kaneda, N., see Nakahara, D., 44
Kasuya, Y., see Kamei, J., 213
Kataoka, K., see Zhang, L., 105
Katoh, Y., see Takeuchi, Y., 71
Kawakami, T., see Kusakabe, T., 59
Kawase, T., see Shimizu, K., 9
Kawashima, N., see Kamei, J., 213
Kewitz, B., see Weiler, R., 151
Kimata, K., see Yamamoto, M., 39
Kobashi, M., Ichikawa, H., Sugimoto, T. and Adachi, A., Response of neurons in the solitary tract nucleus, area postrema and lateral parabrachial nucleus to gastric load of hypertonic saline, 47
Kostakos, M., Persinger, M.A. and Peredery, O., Deficits in working but not reference memory in adult rats in which limbic seizures had been induced before weaning: implications for early brain injuries, 209
Koyanagi, T., see Matsuura, T., 5
Kudoh, T., see Kuwahara, T., 1
Kumar, K.N., see Eggeman, K.T., 173
Kusakabe, T., Kawakami, T., Hori, H., Bandou, Y. and Takenaka, T., Immunohistochemical coexistence of calcitonin gene-related peptide and substance P in the nerve fibers of the internal gills of bullfrog (*Rana catesbeiana*) larvae, 59
Kuwahara, T., Kudoh, T., Nakano, A., Yoshizaki, H., Takamiya, M., Nagase, H. and Arisawa, M., Species specificity of pharmacological characteristics of CCK-B receptors, 1

Łakomy, M., see Całka, J., 21
Legrand, A.M., see Molgó, J., 147
Levine, J.D., see Green, P.G., 117
Li, M., see Yamamoto, M., 39
Li, Y.-Q., Takada, M. and Mizuno, N., Demonstration of habenular neurons which receive afferent fibers from the nucleus accumbens and send their axons to the midbrain periaqueductal gray, 55
Lin, J.-H., see Gean, P.-W., 170
Lisý, V. and Šťastný, F., Effect of repeated hyperammonemia on Na^+-dependent binding of glutamate in rat cortical and hippocampal synaptic membranes, 113
Luboldt, W., see Dudel, J., 177
Luo, J., see Green, P.G., 117

Ma, J., see Aou, S., 197
Mailleux, P., see Go, M., 135
Maitra, K.K., see Seth, P., 217
Majewski, M., see Całka, J., 21
Malcangio, M. and Bowery, N.G., $GABA_B$ receptor-mediated inhibition of forskolin-stimulated cyclic AMP accumulation in rat spinal cord, 189
Martinez, R.P., see Raffa, R.B., 87
Martínez-Murillo, R., see Villalba, R.M., 109
Massa, R., see Sancesario, G., 101
Masuda, S., see Zhang, L., 105
Mataga, N., see Osborne, P.G., 201
Matsuda, T., see Ikeuchi, Y., 63
Matsui, T., see Ohta, A., 33
Matsuura, T., Ogata, A., Demura, T., Moriwaka, F., Tashiro, K., Koyanagi, T. and Nagashima, K., Identification of androgen receptor in the rat spinal motoneurons. Immunohistochemical and immunoblotting analyses with monoclonal antibody, 5
Matyja, E. and Albrecht, J., Ultrastructural evidence that mercuric chloride lowers the threshold for glutamate neurotoxicity in an organotypic culture of rat cerebellum, 155
McKenna, M.M., see Pellis, S.M., 13
McKinney, W.T., see Ginsberg, S.D., 130
Messamore, E., Warpman, U., Williams, E. and Giacobini, E., Muscarinic receptors mediate attenuation of extracellular acetylcholine levels in rat cerebral cortex after cholinesterase inhibition, 205
Michaelis, E.K., see Eggeman, K.T., 173
Mio, T., see Ikeuchi, Y., 63
Misawa, M., see Kamei, J., 213
Mitani, A., see Zhang, L., 105
Mitani, S., see Shimizu, K., 9
Mitsuma, T., see Yamamoto, M., 39
Mizuno, N., see Li, Y.-Q., 55
Mokrushin, A.A. and Emelyanov, N.A., Frequency-dependent plasticity of potentials evoked by repetitive stimulation of the lateral olfactory tract in rat olfactory cortex slices, 16
Molgó, J., Shimahara, T. and Legrand, A.M., Ciguatoxin, extracted from poisonous morays eels, causes sodium-dependent calcium mobilization in NG108-15 neuroblastoma × glioma hybrid cells, 147
Morello, M., see Sancesario, G., 101
Morita, I., see Ikeda, J., 193
Moriwaka, F., see Matsuura, T., 5
Morrison, J.H., see Ginsberg, S.D., 130
Murota, S.-i., see Ikeda, J., 193
Mutoh, T., see Yamamoto, M., 39

Nagase, H., see Kuwahara, T., 1
Nagashima, K., see Matsuura, T., 5
Nagatsu, T., see Nakahara, D., 44
Nakahara, D., Hashiguti, H., Kaneda, N., Sasaoka, T. and Nagatsu, T., Normalization of tyrosine hydroxylase activity in vivo in the striatum of transgenic mice carrying human tyrosine hydroxylase gene: a microdialysis study, 44
Nakajima, Y., see Sakamoto, T., 92
Nakamura, S., see Yamashita, H., 229
Nakano, A., see Kuwahara, T., 1
Negro, M., Casanova, E., Chinchetru, M.A., Fernández-López, A. and Calvo, P., Differential effect of chronic ethanol treatment on barbiturate and steroid modulation of muscimol-binding to rat brain cortex, 83
Nek, N., Schwegler, H., Crusio, W.E. and Frotscher, M., Are the fine-structural characteristics of mouse hippocampal mossy fiber synapses determined by the density of mossy fiber axons?, 75
Nii, Y., see Taniguchi, K., 185

O'Donoghue, D.L., Poff, C.R.D. and Block, J.J., Chronic neonatal *N*-methyl-D-aspartate receptor antagonism with MK-801 increases the number of corticospinal cells retained into adulthood in the rat, 143
Ochiai, K., see Ikeda, J., 193
Ogata, A., see Matsuura, T., 5
Ogawa, K., see Taniguchi, K., 185
Ogino, M., see Shimizu, K., 9

Ohshima, H., see Tanaka, K., 67
Ohta, A., Takagi, H., Matsui, T., Hamai, Y., Iida, S. and Esumi, H., Localization of nitric oxide synthase-immunoreactive neurons in the solitary nucleus and ventrolateral medulla oblongata of the rat: their relation to catecholaminergic neurons, 33
Okada, Y., see Ikeuchi, Y., 63
Onaya, T., see Go, M., 135
Onoe, H., see Osborne, P.G., 201
Osborne, P.G., Mataga, N., Onoe, H. and Watanabe, Y., Behavioral activation by stimulation of a GABAergic mechanism in the preoptic area of rat, 201

Pal, R., see Eggeman, K.T., 173
Pellis, S.M., Castañeda, E., McKenna, M.M., Tran-Nguyen, L. and Whishaw, I.Q., The role of the striatum in organizing sequences of play fighting in neonatally dopamine-depleted rats, 13
Peredery, O., see Kostakos, M., 209
Persinger, M.A., see Kostakos, M., 209
Poff, C.R.D., see O'Donoghue, D.L., 143

Raffa, R.B., Martinez, R.P. and Connelly, C.D., Scyliorhinin-I and -II induce reciprocal hindlimb scratching in mice: differentiation of spinal and supraspinal neurokinin receptors in vivo, 87
Ringler, N.J., see Schengrund, C.-L., 159
Rivera, R., see Ferrer, I., 163
Rodrigo, J., see Villalba, R.M., 109
Ross, H.-G., see Seth, P., 217

Saito, R., see Shimizu, K., 9
Sakamoto, T., Yamanaka, Y., Wada, K. and Nakajima, Y., Effects of tracheostomy on electrically induced vocalization in decerebrate cats, 92
Sakurai, Y., Dependence of functional synaptic connections of hippocampal and neocortical neurons on types of memory, 181
Sancesario, G., Morello, M., Massa, R., Fusco, F. and Bernardi, G., NADPH diaphorase activity is inhibited by EDTA in neurons but not in choroid plexus epithelium, 101
Sasaoka, T., see Nakahara, D., 44
Schaffner, A.E., see Vautrin, J., 125
Schengrund, C.-L., DasGupta, B.R. and Ringler, N.J., Ganglioside GD3 enhances adherence of botulinum and tetanus neurotoxins to bovine brain synapsin I, 159
Schiffmann, S.N. and Vanderhaeghen, J.-J., Age-related loss of mRNA encoding adenosine A_2 receptor in the rat striatum, 121
Schwegler, H., see Nek, N., 75
Seth, P., Gajendiran, M., Maitra, K.K., Ross, H.-G. and Ganguly, D.K., Evidence for D1 dopamine receptor-mediated modulation of the synaptic transmission from motor axon collaterals to Renshaw cells in the rat spinal cord, 217
Shimahara, T., see Molgó, J., 147
Shimizu, K., Ogino, M., Yamaguchi, N., Mitani, S., Saito, R., Kawase, T. and Toya, S., Propentofylline (HWA285) inhibits the release of dopamine during transient ischemia and modulates its metabolism in rat striatum, 9
Sobue, G., see Yamamoto, M., 39
Solano-Flores, L.P., see Hochstenbach, S.L., 225
Song, Z.-M., Brookes, S.J.H. and Costa, M., NADPH-diaphorase reactivity in nerves supplying the rat anococcygeus muscle, 221
Soria, M., see Ferrer, I., 163
Spreafico, R., De Biasi, S., Amadeo, A. and De Blas, A.L., $GABA_A$-receptor immunoreactivity in the rat dorsal thalamus: an ultrastructural investigation, 232
Šťastný, F., see Lisý, V., 113
Stoof, J.C., see Henselmans, J.M.L., 79
Sugimoto, T., see Kobashi, M., 47
Suzuki, T., see Kamei, J., 213

Takada, M., see Li, Y.-Q., 55
Takagi, H., see Ohta, A., 33
Takahashi, K., see Takeuchi, Y., 71
Takamiya, M., see Kuwahara, T., 1
Takazawa, K., see Go, M., 135
Takenaka, T., see Kusakabe, T., 59
Takeuchi, Y., Katoh, Y. and Takahashi, K., Classical acetylcholine receptors do not play a direct role in neuronal transmission of photic information in the suprachiasmatic nucleus in rats, 71
Tanaka, K., Ohshima, H., Esumi, H. and Chiba, T., Direct synaptic contacts of nitric oxide synthase-immunoreactive nerve terminals on the neurons of the intracardiac ganglia of the guinea pig, 67
Taniguchi, K., Nii, Y. and Ogawa, K., Subdivisions of the accessory olfactory bulb, as demonstrated by lectin-histochemistry in the golden hamster, 185
Tao, Y., see Droge, M.H., 139
Tashiro, K., see Matsuura, T., 5
Toda, T. and Hayashi, H., The inhibitory effect of substance P antagonist, CP-96,345, on the late discharges of nociceptive neurons in the rat superficial spinal dorsal horn, 36
Toya, S., see Shimizu, K., 9
Tran-Nguyen, L., see Pellis, S.M., 13

Uchida, T., see Go, M., 135

Vanderhaeghen, J.-J., see Schiffmann, S.N., 121
Vautrin, J., Schaffner, A.E. and Barker, J.L., Tonic GABA secretion of cultured rat hippocampal neurons rapidly transformed by Zn^{2+} into quantal release, 125
Vidal, A., see Ferrer, I., 163
Villalba, R.M., Rodrigo, J. and Martínez-Murillo, R., Presence of C-flanking peptide of neuropeptide Y (C-PON)-immunoreactive neurons in the olfactory cortex of the hedgehog (*Erinaceus europaeus*), 109

Wada, K., see Sakamoto, T., 92
Walsh, J., see Eggeman, K.T., 173
Warpman, U., see Messamore, E., 205
Watanabe, Y., see Osborne, P.G., 201
Weiler, R. and Kewitz, B., The marker for nitric oxide synthase, NADPH-diaphorase, co-localizes with GABA in horizontal cells and cells of the inner retina in the carp retina, 151
Whishaw, I.Q., see Pellis, S.M., 13
White, T.D., see Craig, C.G., 167
Williams, E., see Messamore, E., 205
Wu, S.Y., see Dun, N.J., 51

Yamaguchi, N., see Shimizu, K., 9
Yamamoto, M., Sobue, G., Mutoh, T., Li, M., Doyu, M., Mitsuma, T. and Kimata, K., Gene expression of high- ($p140^{trk}$) and low-affinity nerve growth factor receptor (LNGFR) in the adult and aged human peripheral nervous system, 39
Yamanaka, Y., see Sakamoto, T., 92
Yamashita, H., Zhang, Y.-X. and Nakamura, S., The effects of thiamin and its phosphate esters on dopamine release in the rat striatum, 229
Yoshizaki, H., see Kuwahara, T., 1

Zhang, L., Andou, Y., Masuda, S., Mitani, A. and Kataoka, K., Dantrolene protects against ischemic, delayed neuronal death in gerbil brain, 105
Zhang, Y.-X., see Yamashita, H., 229
Zújar, M.J., see Ferrer, I., 163